Pre-	Operative Regimen	SURGERY DATE:	/ /							
	,	RIGHT EYE	LEFT EYE							
員	PROLENSA / ILEVRO: 1 drop 1x/day, start 2 days before surgery									
□ PROLENSA / ILEVRO: 1 drop 1x/day, start 2 days before surgon □ BROMSITE: 1 drop 2x/day, start 2 days before surgery										
	□ PRN Dry Eye Omega Benefits / Hydroeyes <sup>®</sup>									
TWO	<b>Liquid:</b> 1 teaspoon	iquid: 1 teaspoon per day								
	Softgels: 4 per day	Softgels: 4 per day								
u d	Start Omega 3s as									
	☐ HYPOCHLOR/ AVE	NOVA / OCUSOFT ®:	MC(MT CR T							
	in School Ave	NAC (NATI CIETS								

GANII

Use lid cleaner daily or until finished. Start 3 days prior to surgery in both eyes

☐ FRESHKOTE® PF:

1 drop in both eyes 3 or 4 times daily for 1 month



### **General Instructions**

- Follow all your primary care physician's instructions regarding all medications.
- Notify Dr. Cohen if you are taking Flomax or prostate medications.
- Take medications only as directed with a small sip of water.
- Wash hands before using drops.
- Wait 5 MINUTES between different drops.
- If you currently take other eye drops, please continue to take them as you normally do unless instructed otherwise.

# Day of Surgery Regimen

- Use one drop of PROLENSA or BROMSITE when you wake up the morning of surgery.
- Do not eat or drink anything after midnight the night before surgery.
- After surgery, take your daily dose of PRN Dry Eye Omega Benefits® or HydroEye®

ANTIBRIOTIC

#### **PREVENTS INFECTION**

- ☐ **BESIVANCE:** One drop 3x/day for 1-2 weeks (as directed)
- ☐ **TOBRA/POLYMYXIN:** One drop 4x/day for 1-2 weeks (as directed)
- ☐ MOXI/CIPROFLOX: One drop 4x/day for 1-2 weeks



### PREVENTS INFLAMMATION AND PAIN

- ☐ **BROMSITE:** One drop 2x/day for 1 week, then 1x/day for 3 weeks
- ☐ **PROLENSA:** One drop 1x/day for 4-6 weeks (as directed)
- ☐ **ILEVRO:** One drop 1x/day for 4-6 weeks (as directed)




Lotemax

STEROID

#### REGULATES HEALING AND INFLAMMATION

- ☐ TOBRADEX ST\* / FLAREX 0.1%: One drop 4x/day for 3-6 weeks taper (as directed)
- □ LOTEMAX SM / PREDNISOLONE: One drop 3-4x/day for 4-6 weeks taper (as directed)



DROP START	PRE-OP	DAY OF	WEEK 1	WEEK 2	WEEK 3	WEEK 4
DATE <del>-&gt;</del>	is.	c				
ANTIBIOTIC		2x	4x	stop		
NSAID	1-2x	2x	1 - 2x	1x	1x	0 - 1x
STEROID*		2x	2 - 4x	2 - 3x	1 - 2x	0 - 1x

### **Post-Operative Reminders**

- DO NOT RUB THE OPERATED EYE.
- Wear eye shield at night for 1 week.
- Avoid heavy lifting, bending, straining, and exercising for 1 week.
- Wear your old glasses if more comfortable. **Wear sunglasses** or tinted shades while outside during the day if more comfortable.
- You may take a full-body shower and shampoo 24 hrs after surgery. Close eyes to avoid getting water in operated eye.
- Avoid swimming and gardening for 2 weeks.
- Do not wear eye makeup for 1 week.
- Most people can resume driving several days after surgery.
  Use your best judgment.
- Bring all medications to our office on each post-operative visit.
- Continue taking **PRN** or **Hydroeye Omega Benefits** and **FRESHKOTE PF** \* post-surgery to make sure your eye is producing a healthy tear and to improve quality of vision.
- Tylenol may be taken for discomfort. No Advil or aspirin unless approved.

### **Normal Post-Operative Symptoms**

**Blurred** vision, **halos** around light, eye **tearing**, eye **redness**, and small amount of **mucous** that collects on the eyelids is normal after surgery. A slight **burning** sensation from the drops is also expected. Symptoms should **improve gradually**.

## THE FOLLOWING SYMPTOMS REQUIRE URGENT ATTENTION:

- Extreme pain or swelling
- Significant discharge or pus
- Sudden vision loss
- Flashing lights or floaters



**Cohen Laser and Vision Center** 

3020 N Military Trail, Suite 150 Boca Raton, FL 33431 P (561) 981-8400 www.cohenlaser.com